

# Chapter 1

## Importance of Carp Genetic Resources<sup>1</sup>

Modadugu V. Gupta<sup>2</sup>, Madan M. Dey<sup>2</sup> and David J. Penman<sup>3</sup>

Fish and fisheries play an important role in the economies of developing countries, contributing to animal protein intake, employment generation, household incomes and foreign exchange earnings. Surveys conducted by the WorldFish Center and FAO show that fish has become an increasingly important source of protein over the last decade in most of the developing countries. Countries with low per capita gross domestic product tend to have a higher share of fish protein in their animal protein consumption (Kent 1997). Studies indicated that demand for fish increases as expenditure/income rises and that higher income groups tend to consume more fish than lower income groups. However, the share of fish (as protein) and the share of fish to the total food expenditure are higher among lower income groups, suggesting that lower income groups are the most dependent on fish (Dey et al. 2004; FAO 1999a; Dey 2000; Dey et al. 2000a; ICLARM 2000). This result is also consistent with the generalization that although less developed countries are not the biggest consumers of fish, they are the most dependent on aquatic resources (FAO 1993; Kent 1997; FAO 1999b), indicating the importance of fish as a primary source of protein among relatively poorer households in these countries. No wonder it is regarded as “poor man’s protein” (Williams 1996).

In Asia, on average, almost 30 per cent of the total animal protein intake is derived from fish. Among the Southeast Asian countries, fish protein provides 45 per cent of the total protein consumed (Prein and Ahmed 2000). Although Japan, the European Union and USA have higher per capita consumption of fish and fish products, the share of fish protein in terms of the total animal protein consumption is far less than that in many developing countries.

World fish production in 1999 was estimated at 130 million t, of which only 97 million t was for

human consumption (FAO 2003). FAO (1999a) estimated that by the year 2010, demand for food fish will have increased by 13.5-18 per cent or to about 105-110 million t. Against this demand, production from capture fisheries is declining as most of the world’s fish stocks have been exploited to their maximum potential or over exploited, indicating it might be difficult to increase yields from capture fisheries in the near future (Williams 1996). During the 1950s and 1960s capture fish production increased by an average of 6 per cent per annum and this declined to 2 per cent during the 1970s and 1980s, falling almost to zero during the 1990s (FAO 2000).

In contrast to declining growth in capture fisheries, aquaculture has been growing at a fast rate, by about 5 per cent per year during the 1950s and 1960s to about 8 per cent during the 1970s and 1980s and over 10 per cent since 1990 (FAO 2000). Aquaculture production has increased from 3.2 per cent of the total fish production in 1950 to 32 per cent in 2000. Aquaculture has become the world’s fastest growing food-producing sector, with production more than doubling during 1990-2000, from 13.1 million t in 1990 to 35.6 million t in 2000 (FAO 2003).

Of the total aquaculture production of 45.7 million t in 2000, 45 per cent was contributed by freshwater aquaculture. Production increased from 7.9 million t in 1991 to 20.6 million t in 2000. Asia produces about 91 per cent of the world’s freshwater aquaculture production, with China contributing 74 per cent of this (Fig.1). Other major producers of freshwater aquaculture are India (9.89 per cent), Bangladesh (2.77 per cent), Vietnam (1.85 per cent), Indonesia (1.76 per cent), Thailand (1.29 per cent) and the Philippines (0.54 per cent). Growth in freshwater aquaculture in the last decade among these countries differs: Vietnam had the highest growth

<sup>1</sup> WorldFish Center Contribution No. 1728.

<sup>2</sup> WorldFish Center, Jalan Batu Maung, Batu Maung, 11960 Bayan Lepas, Penang, Malaysia.

<sup>3</sup> University of Stirling, U.K.

of 14.54 per cent per annum during 1990-2000, followed by China (13.6 per cent), Bangladesh (12.94 per cent), Thailand (9.38 per cent) and Indonesia (5.65 per cent).

With this remarkable increase in production, relatively low prices and limited international markets, freshwater fish are expected to become an increasingly important source of animal protein, particularly for those in medium and lower income groups in developing countries, especially in Asia (Dey et al. 2004).


About 242 species of freshwater finfish are cultured globally and of these, carps (Cyprinidae) are the dominant species, especially in Asia. While production of cultured carps is on the increase the amount from capture fisheries from rivers and other natural waters is on the decline due to the shrinking of areas for spawning, over fishing and other human induced habitat degradation. Production of carps from culture has increased from 5.6 million t in 1990 to 16.4 million t in 2001, while the yield from capture fisheries has declined from 0.75 million t in 1990 to 0.54 million t in 2001 (Table 1.1). These human induced changes are leading to threats to the existence of some of the species (Acosta and Gupta, Chapter 7, this vol.). In Asia, aquaculture

production increased from 10.8 million t in 1990 to 29.48 million t in 2001, while carp production increased from 5.07 million t in 1990 to 16.42 million t in 2001 (Fig. 1.1).

**Table 1.1. Carp production (million t) from capture fisheries and culture during 1990-2001**

Year	Production from culture (million t)	Production from capture fisheries (million t)	Total (million t)
1990	5.07	0.47	5.54
1991	5.27	0.54	5.81
1992	6.08	0.39	6.47
1993	7.18	0.45	7.63
1994	8.50	0.51	9.01
1995	9.93	0.55	10.48
1996	11.48	0.43	11.90
1997	12.94	0.38	13.32
1998	13.60	0.38	13.98
1999	14.63	0.38	15.01
2000	15.14	0.33	15.47
2001	16.00	0.31	16.31
Growth rate (%)	11.16%	-2.60	10.44%

Source: FAO 2003. Fisheries Statistics (on line) available at [www.fao.org/fi/statist/FISOFT/FISHPLUS.asp](http://www.fao.org/fi/statist/FISOFT/FISHPLUS.asp) [June 2003].



**Fig. 1.1. Trends in world aquaculture and carp production (million t) from aquaculture globally and from Asia**

**Table 1.2. Prominent Cyprinid species used for aquaculture in Asia**

Species	Production (t) in 2001 <sup>a</sup>
<i>Hypophthalmichthys molitrix</i>	3 546 285
<i>Ctenopharyngodon idella</i>	3 636 367
<i>Cyprinus carpio</i>	2 849 492
<i>Aristichthys nobilis</i>	1 663 499
<i>Carassius carassius</i>	1 527 058
<i>Labeo rohita</i>	833 816
<i>Catla catla</i>	870 085
<i>Cirrhinus cirrhosus</i>	589 841
<i>Parabramis pekinensis</i>	541 115
<i>Mylopharyngodon piceus</i>	190 707
<i>Cirrhinus molitorella</i>	220 118
<i>Barbonymus gonionotus<sup>b</sup></i>	53 563
<i>Puntius javanicus<sup>b</sup></i>	26 251
<i>Osteochilus hasseltii</i>	15 319
<i>Leptobarbus hoevenii</i>	1 111
Other cyprinid species	56 440
TOTAL	16 621 067

<sup>a</sup> Represents production of carp from aquaculture in 23 countries of Asia.


<sup>b</sup> *Puntius gonionotus* and *Puntius javanicus* are the same species, now described as *Barbonymus (Barbodes) gonionotus* by FishBase (2003) [available on line at: www.fishbase.org].

Source: FAO 2003. Fishery statistics: Aquaculture production. Vol. 92/2

Over 99 per cent of the total carp production was contributed by 15 species among which Chinese carps [silver carp (*Hypophthalmichthys molitrix*), grass carp (*Ctenopharyngodon idella*), bighead carp (*Aristichthys nobilis*), crucian carp (*Carassius carassius*) and common carp (*Cyprinus carpio*)] and Indian carps [rohu (*Labeo rohita*), catla (*Catla catla*) and mrigal (*Cirrhinus cirrhosus*)] are the dominant species (Table 1.2 and Fig. 1.2).

Although aquaculture has a long history in Asia, dating back over 2000 years in China, until recently aquaculture has not been subject to the same intensive development of its genetic base as has been the case with terrestrial agriculture. Genetic improvement of fish is estimated to lag behind advances in livestock by nearly 50 years (Eknath et al. 1991). In fact until the 1960s, the aquaculture industry was dependent on seed of carps collected from the wild, but with successful hypophysation of carps in the late 1950s, much of the seed is presently being produced in hatcheries. With this development, there is increasing concern that the genetic bases of many culture species are deteriorating to the extent of lowering the growth (Eknath and Doyle 1990) and the most common stocks used in aquaculture may not be the best performers (e.g. Nile tilapia in the Philippines, Eknath et al. 1993; Dey et al. 2000b; Dey and Gupta 2000). Evidence of inbreeding and unintentional selection among hatchery-based and established farm stocks have been observed (Pullin and Capilli 1988; McAndrew et al. 1993; Eknath 1995).

While remarkable progress has been made in improving the productivity of crops and livestock in the last three to four decades through breeding and selection, it is only in recent years that efforts have been made to harness the benefits of genetic enhancement (e.g. salmon and Nile tilapia). Selective breeding of Nile tilapia (*Oreochromis niloticus*) over five generations has resulted in an



**Fig. 1.2. Percentage of contribution of major carp species to the total carp production from aquaculture in Asia during 2001**

improved strain, the GIFT strain that has shown 85 per cent higher growth as compared to the existing stocks in the Philippines (Eknath and Acosta 1998). Furthermore, testing of the GIFT strain through on-station and on-farm trials indicated the GIFT strain's faster growth as compared to the existing strains in Bangladesh, China, Thailand and Vietnam (Dey et al. 2000b). This has created interest among a number of major aquaculture producing countries in Asia and research is in progress for genetic improvement of *C. carpio*, *L. rohita*, silver barb (*B. gonionotus*) and Blunt snout bream (*Megalobrama amblycephala*), besides others, with funding support from the Asian Development Bank (WorldFish Center 2001). While most of these genetic improvement programs have as yet had little impact on aquaculture, there are some examples that demonstrate the kind of impact that can hopefully be expected to become much more widespread in the near future. Over one million tonnes of the Jian strain of common carp is produced each year in China, and several other improved varieties are cultured on a significant scale in China. Genetically improved varieties of several other cyprinid species have been developed and in some cases disseminated in several Asian countries.

In view of the significant growth of aquaculture and the need for increased production, there is a tendency among farmers and government agencies to introduce exotic species. The available records indicate that 259 introductions comprising 42 cyprinid species have taken place in 27 countries of Asia (Acosta and Gupta, Chapter 7, this vol.). In most cases these new species have led to improved aquaculture production, while in some cases they have also resulted in adverse impacts on the aquatic ecosystem.

Conservation and proper utilization of germplasm is a prerequisite for the sustainable management of and increased production from aquatic resources. In view of this, several major carp producing countries in Asia (Bangladesh, China, India, Indonesia, Thailand and Vietnam) have been involved in genetic improvement of carps under a project funded by the Asian Development Bank. As part of this, a review of the carp genetic resources in these countries was undertaken. The results of this are detailed in Chapter 3. The publication also includes a review of progress of genetics research in carps, introductions of carps and their impacts.

## References

- Dey, M.M. 2000. Analysis of demand for fish in Bangladesh. *Aquaculture Economics and Management* 4(1&2):63-82.
- Dey, M.M. and M.V. Gupta. 2000. Socioeconomic aspects of disseminating genetically improved Nile tilapia (*Oreochromis niloticus*) in Asia: Introduction. *Aquaculture Economics and Management* 4(1&2): 5-12.
- Dey, M.M., G.P. Bimbao, L. Yong, P. Regaspi, A.H.M. Kohinoor, Do Kim Chung, N. Pongthana and F.J. Paraguas. 2000a. Current status of production and consumption of tilapia in selected Asian countries. *Aquaculture Economics and Management* 4(1&2): 47-62.
- Dey, M.M., A.E. Eknath, Li Sifa, M.G. Hussain, T.M. Tran, M. Pongthana, V.H. Nguyen and F.J. Paraguas. 2000b. Performance and nature of genetically improved farmed tilapia: A bioeconomic analysis. *Aquaculture Economics and Management* 4(1&2):83-106.
- Dey, M.M., M.A. Rab, F.J. Paraguas, S. Piumsombun, R. Bhatta, Md. F. Alam, W. Miao and M. Ahmed. 2004. Fish consumption in selected Asian countries: A disaggregated analysis by types of fish and classes of consumers. *Aquaculture Economics and Management* (In press).
- Eknath, A.E. 1995. Managing aquatic genetic resources. Management example 4: The Nile tilapia, p. 176-194. *In* J.Thorpe, G. Gall, J.E. Lannan and C.E. Nash (eds). *Conservation of fish and shellfish resources: managing diversity*. Academic Press, Harcourt Brace Company, London.
- Eknath, A.E and B.O. Acosta. 1998. Genetic improvement of farmed tilapias (GIFT) project. Final Report, March 1988 to December 1997. International Center for Living Aquatic Resources Management, Manila, Philippines (mimeo).
- Eknath, A.E. and R.W. Doyle. 1990. Effective population size and rate of inbreeding in aquaculture of Indian major carps. *Aquaculture* 85 (1/4):293-305.
- Eknath, A.E., H.B. Bentsen, B. Gjerde, M.M. Tayamen, T.A. Abella, T. Gjedrem and R.S.V. Pullin. 1991. Approaches to national fish breeding programs: Pointers from a tilapia pilot study. *NAGA, the ICLARM Quarterly* 14(2): 10-12.
- Eknath, A.E., M.M. Tayamen, M.S. Palada de Vera, J.C. Danting, R.A. Reyes, E.E. Dionisio, J.B. Capilli, H.L. Bolivar, T.A. Abella, A.V. Circa, H.B. Bentsen, B. Gjerde, T. Gjedrem and R.S.V. Pullin. 1993. Genetic improvement of farmed tilapias: The growth performance of eight strains of *Oreochromis niloticus* tested in different farm environments. *Aquaculture* 111 (1/4):171-188.
- FAO. 1993. Availability of fish supplies for international trade. *FAO COFI:FT/IV/93/3*.
- FAO. 1999a. Fish and Fishery Products: World apparent consumption statistics based on food balance sheet. *FAO Fisheries Circular*, No. 821 (Rev. 5).
- FAO. 1999b. The State of World Fisheries and Aquaculture. *FAO, Rome*.

- FAO. 2000. The state of world fisheries and aquaculture. FAO, Rome.
- FAO. 2003. Fisheries statistics – aquaculture production. 2001. Available online at <http://www.fao.org/fi/statist/FISOFT/FISHPLUS.asp> [June 2003].
- ICLARM. 2000. Genetic Improvement of Carp Species in Asia: Final Report. Asian Development Bank Regional Technical Assistance No. 5711. ICLARM, Penang, Malaysia.
- Kent, G. 1997. Fisheries, food security and the poor. *Food Policy* 22(5):393-404.
- McAndrew, B.J., K.J. Rana and D.J. Penman. 1993. Conservation and preservation of genetic variation in aquatic organisms, p. 295-336. *In* J.F. Muir and R.J. Roberts (eds). *Recent Developments in Aquaculture IV*. Blackwell Scientific Publications, Oxford, U.K.
- Prein, M. and M. Ahmed. 2000. Integration of aquaculture into smallholder farming systems for improved food security and household nutrition. *Food and Nutrition Bulletin*, 21(4):466-471.
- Pullin, R.S.V. and J.B. Capilli. 1988. Genetic improvement of tilapias: problems and prospects, p. 259-266. *In* R.S.V. Pullin, T. Bhukasawan, K. Tonguthai and J.L. Maclean (eds). *The Second International Symposium on Tilapia in Aquaculture*. ICLARM Conf. Proc. 15, 623 p.
- Williams, M. 1996. The transition in the contribution of living aquatic resources to food security. IFPRI Food, Agriculture and the Environment Discussion Paper, No. 13. 41 p.
- WorldFish Center. 2001. Final Report of the Genetic Improvement of Carps project. ICLARM-WorldFish Center (mimeo).